



Residential Aged Care (RAC) Resources

palliaged.com.au

palliAGED is funded by the Australian Government Department of Health and Aged Care and managed by CareSearch, Flinders University

palliAGED

The online palliative care evidence and practice information resource for the aged care sector

To obtain further copies of this pack:

- Download or order printed copies at palliaged.com.au/practiceforms



Find out more:

- Go to palliaged.com.au
- Email us at palliaged@flinders.edu.au
- Follow us on  Twitter [@palliaged](https://twitter.com/palliaged)
- Sign up to receive our newsletter and articles at palliaged.com.au/subscribe





Residential Aged Care Pack Contents

Self-Care Plan - Aged Care Team

To care for others, you need to take care of yourself. This worksheet helps you to plan activities to maintain your balance in life.

- Self-care plan for the Aged Care Team.

Recognising changing needs - forms to help you identify changes

If you see clients regularly, it can be more difficult to notice subtle changes in their condition. These forms can be used to systematically assess for deterioration in a person's health and to identify unmet supportive and palliative care needs.

- SPICT Tool
- SPICT4ALL Tool.

Symptom control

Pain is one of the most common symptoms in palliative care. You can use this form to monitor for changes in pain status.

- Abbey Pain scale.

Organising a palliative care case conference

A series of practical checklists and forms to guide and document case conferences.

- Using the palliAGED conference forms
- Case Conference Checklist - residential care
- Invitation to attend a case conference - GPs
- Confirmation of a case conference - GPs
- Information about case conferences and palliative care
- Invitation to attend a case conference - person and family
- Confirmation for you and your family
- Staff Communication Form
- Case Conference Summary - residential care.

End of life care

Practical lists of medicines that can be used for responding to unanticipated needs in the terminal phase.

- Medicines from the PBS prescriber's bag for terminal phase symptoms.

MBS Remuneration

- MBS items for nurse practitioners
- MBS Remuneration to Support a Planned General Practice Palliative Care Pathway (Residential Aged Care).

To obtain further copies of this pack:

Download or order printed copies at palliaged.com.au/practiceforms



palliaged.com.au

palliAGED is funded by the Australian Government Department of Health and Aged Care and managed by CareSearch, Flinders University

This page is intentionally left blank

Self-Care Plan:

Aged Care Team

Name:

Caring for others can be rewarding. However, staff working in aged care look after many people who die. You might find this loss hard to accept. You might find it hard to sleep, no longer enjoy your work, or feel tired. This can affect you and your family. Self-care is what we do to maintain balance in our life.

A self-care plan based on what you like to do can help. We have suggested a few things that you could try, but what you choose will depend on you.

Workplace self-care - Activities to help you at work

Some examples

Regular meetings with supervisors or a more experienced colleague

Join a support group with the people you work with

Attend training programs

My Activities:

Physical self-care - Activities that help you to stay fit and healthy

Some examples

Develop a regular sleep routine

Aim for a healthy diet

Take lunch breaks and go for a walk

Get some exercise before/after work regularly

My Activities:

Self-Care Plan (continued)

Psychological self-care - Activities that help you to feel clear-headed and able to engage with workplace and personal challenges

Some examples

Keep a reflective journal

Seek regular meetings with supervisors or a more experienced colleague

Turn off your email and work phone outside of work hours

Make time to be with friends and family

My Activities:

Emotional self-care - Allowing yourself to safely express your emotions

Some examples

Develop friendships that are supportive

Write or think of three good things that you did each day

Play a sport and have a coffee together after training

Talk to your friends about how you are coping with work and life demands

My Activities:

Adapted with permission from ReachOut Australia

Self-Care Plan (continued)

Spiritual self-care - Develop a sense of perspective beyond the day-to-day of life which can include religion, but it is not always about religion

Some examples

Engage in reflective practices like meditation

Go on walks to connect with nature

Go to church/mosque/temple

Do yoga

My Activities:

Relationship self-care - Maintain healthy, supportive relationships, and ensure that you are not only connected to work people

Some examples

Prioritise close relationships in your life eg, with partners, family and children

Attend the special events of your family and friends

Arrive to work and leave on time every day

My Activities:

Adapted with permission from ReachOut Australia

This page is intentionally left blank



Supportive and Palliative Care Indicators Tool (SPICT™)

The SPICT™ is used to help identify people whose health is deteriorating. Assess them for unmet supportive and palliative care needs. Plan care.

Look for any general indicators of poor or deteriorating health.

- Unplanned hospital admission(s).
- Performance status is poor or deteriorating, with limited reversibility. (eg. The person stays in bed or in a chair for more than half the day.)
- Depends on others for care due to increasing physical and/or mental health problems.
- The person's carer needs more help and support.
- Progressive weight loss; remains underweight; low muscle mass.
- Persistent symptoms despite optimal treatment of underlying condition(s).
- The person (or family) asks for palliative care; chooses to reduce, stop or not have treatment; or wishes to focus on quality of life.

Look for clinical indicators of one or multiple life-limiting conditions.

Cancer

Functional ability deteriorating due to progressive cancer.

Too frail for cancer treatment or treatment is for symptom control.

Dementia/ frailty

Unable to dress, walk or eat without help.

Eating and drinking less; difficulty with swallowing.

Urinary and faecal incontinence.

Not able to communicate by speaking; little social interaction.

Frequent falls; fractured femur.

Recurrent febrile episodes or infections; aspiration pneumonia.

Neurological disease

Progressive deterioration in physical and/or cognitive function despite optimal therapy.

Speech problems with increasing difficulty communicating and/or progressive difficulty with swallowing.

Recurrent aspiration pneumonia; breathless or respiratory failure.

Persistent paralysis after stroke with significant loss of function and ongoing disability.

Heart/ vascular disease

Heart failure or extensive, untreatable coronary artery disease; with breathlessness or chest pain at rest or on minimal effort.

Severe, inoperable peripheral vascular disease.

Respiratory disease

Severe, chronic lung disease; with breathlessness at rest or on minimal effort between exacerbations.

Persistent hypoxia needing long term oxygen therapy.

Has needed ventilation for respiratory failure or ventilation is contraindicated.

Other conditions

Deteriorating with other conditions, multiple conditions and/or complications that are not reversible; any treatment available will have a poor outcome.

Kidney disease

Stage 4 or 5 chronic kidney disease (eGFR < 30ml/min) with deteriorating health.

Kidney failure complicating other life limiting conditions or treatments.

Stopping or not starting dialysis.

Liver disease

Cirrhosis with one or more complications in the past year:

- diuretic resistant ascites
- hepatic encephalopathy
- hepatorenal syndrome
- bacterial peritonitis
- recurrent variceal bleeds

Liver transplant is not possible.

Review current care and care planning.

- Review current treatment and medication to ensure the person receives optimal care; minimise polypharmacy.
- Consider referral for specialist assessment if symptoms or problems are complex and difficult to manage.
- Agree a current and future care plan with the person and their family. Support carers.
- Plan ahead early if loss of decision-making capacity is likely.
- Record, share, and review care plans.

This page is intentionally left blank



The SPICT™ helps us to look for people who are less well with one or more health problems. These people need more help and care now, and a plan for care in the future. Ask these questions:

Does this person have signs of poor health or health problems that are getting worse?

- Unplanned (emergency) admission(s) to hospital.
- General health is poor or getting worse; the person never quite recovers from being more unwell. (This means the person is less able to manage day to day life and often stays in bed or in a chair for more than half the day).
- Needs help from others for care due to increasing physical and/ or mental health problems.
- The person's carer needs more help and support.
- Has clearly lost weight over the last few months; or stays too thin.
- Has troublesome symptoms most of the time despite good treatment of their health problems.
- The person (or family) asks for palliative care; chooses to reduce, stop or not have treatment; or wishes to focus on quality of life.

Does this person have any of these health problems?

Cancer

Less able to manage usual activities; health getting poorer.

Not well enough for cancer treatment or treatment is to help with symptoms.

Dementia/frailty

Unable to dress, walk or eat without help.

Eating and drinking less; difficulty with swallowing.

Has lost control of bladder and bowels.

Not able to communicate by speaking; not responding much to other people.

Frequent falls; fractured hip.

Frequent infections; pneumonia.

Nervous system problems

(eg Parkinson's disease, MS, stroke, motor neurone disease)

Physical and mental health are getting worse.

More problems with speaking and communicating; swallowing is getting worse.

Chest infections or pneumonia; breathing problems.

Severe stroke with loss of movement and ongoing disability.

Heart or circulation problems

Heart failure or has had attacks of chest pain. Short of breath when resting, moving or walking a few steps.

Very poor circulation in the legs; surgery is not possible.

Lung problems

Unwell with long term lung problems. Short of breath when resting, moving or walking a few steps even when the chest is at its best.

Needs to use oxygen for most of the day and night.

Has needed treatment with a breathing machine in the hospital.

Other conditions

People who are less well and may die from other health problems or complications. There is no treatment available or it will not work well.

Kidney problems

Kidneys not working well; general health is getting poorer.

Stopping kidney dialysis or choosing supportive care instead of starting dialysis.

Liver problems

Worsening liver problems in the past year with complications like:

- fluid building up in the belly
- being confused at times
- kidneys not working well
- infections
- bleeding from the gullet

A liver transplant is not possible.

What we can do to help this person and their family.

- Start talking with the person and their family about any help needed now and why making plans for care is important in case things change.
- Ask for help and advice from a nurse, doctor or other professional who can assess the person and their family and help plan care.
- We can look at the person's medicines and other treatments to make sure we are giving them the best care or get advice from a specialist if problems are complicated or hard to manage.
- We need to plan early if the person might not be able to decide things in the future.
- We make a record of the care plan and share it with people who need to see it.

This page is intentionally left blank

Abbey Pain Scale

In the following, for each of the following six areas enter pain scores:

Absent = 0; Mild = 1; Moderate = 2; Severe = 3

Patient details
Surname.....
Title.....
Given names.....
DOB..... MRN.....
Address.....
Suburb.....
Postcode.....

Enter Date:					
Enter time:					
Sign entry					
1. Vocalisation e.g., whimpering, groaning, crying.					
2. Facial Expression e.g., looking tense, frowning, grimacing, looking frightened.					
3. Change in body language e.g., fidgeting, rocking, guarding part of body, withdrawn.					
4. Behavioural change e.g., increased confusion, refusing to eat, alteration in usual patterns.					
5. Physiological change e.g., temperature, pulse or blood pressure outside normal limits, perspiring, flushing or pallor.					
6. Physical changes e.g., skin tears, pressure areas, arthritis, contractures, previous injuries.					
Total scores					
Circle the range that matches the total pain score					
0-2 No pain	No pain	No pain	No pain	No pain	No pain
3-7 mild	Mild	Mild	Mild	Mild	Mild
8-13 moderate	Moderate	Moderate	Moderate	Moderate	Moderate
14+ severe	Severe	Severe	Severe	Severe	Severe

Tick the box which matches the type of pain: Acute Chronic Acute on chronic

About Abbey Pain Scale

Purpose: Developed to detect pain in elderly residents with dementia and inability to communicate verbally. It is a 6-item 3 point scale tool.

Description: The Abbey Pain Scale was developed for use in aged care and dementia care. The tool is best used as part of an overall pain management plan. As the tool does not differentiate between distress and pain measuring the effectiveness of any interventions is essential. Use the form to collate recordings across an extended period to facilitate monitoring of responses. The Australian Pain Society recommends using the tool as a movement-based assessment and conducting a **second evaluation one hour after any intervention taken**. Repeat hourly until a score of mild pain is reached and then 4 hourly for 24 hours with treatment for pain as required. Contact the GP or pain team if there is no improvement.

Acknowledgement: Abbey J, et al. The Abbey pain scale: a 1-minute numerical indicator for people with end-stage dementia. Int J Palliat Nurs. 2004 Jan;10(1):6-13.

Using the palliAGED Palliative Care Case Conference forms

A case conference or family meeting between the person, their family and care providers can help to explain what is happening and to plan care. The palliAGED forms can help.

1

Use the palliAGED [Case Conference Checklist for residential care](#) or [Case Conference Checklist for home care](#) to organise a palliative care case conference. Tick off items as they are completed.

2

Speak with the person and their family about the need for a case conference. Provide [Information on palliative care and case conferences](#).

3

Involving the person's GP is important. Use the [GP invitation](#) to invite them to attend, and/or to suggest a suitable time.

4

Closer to the date of the Case Conference send a letter [confirming details to the person and their family](#), and send [confirmation to the GP](#).

5

To guide the meeting and to make sure that all steps following the conference are completed use the palliAGED [Case Conference Summary for residential care](#) or [Case Conference Summary for home care sheet](#).

This page is intentionally left blank

Palliative Care Case Conference

Organisation: _____

Planning Checklist - Residential Care

Full name of resident: _____ DOB (dd/mm/yy): _____

Date of case conference (dd/mm/yy): _____ Time: _____

Venue: _____ Room booked: _____

Dial-in telephone number: _____ Code: _____

Case conference facilitator: _____

Goals of case conference:

Family Participants

Name	Role/Relationship	Contact Details

Health and Care Professionals

Name	Role/Relationship	Contact Details

Document (tick as appropriate)	Sent	Accepted/Declined	N/A
Resident & family information		A D	
Resident & family confirmation		A D	
GP invitation		A D	
GP confirmation		A D	

	Needed	Obtained	N/A
Clinical record (including most recent medication chart)			
Advance care planning document (legal or non-legal)			
Other (specify)			

This page is intentionally left blank

GP Invitation

Organisation: _____

Palliative Care Case Conference

To:	Email/fax number:
From:	No. of pages: (including this page)
Subject: Palliative Case Conference	Date sent: (dd/mm/yy):

Dear Dr, _____

A Palliative Care Case Conference is being organized for (Resident/client name):

Resident/client DOB (dd/mm/yy): _____

Proposed date (dd/mm/yy): _____ Start time: _____

Expected duration: _____ Venue: _____

As an important member of the care team for our resident/client, you are invited to participate.

Reason for case conference:

Please indicate availability to participate in this case conference by ticking one of the options below:

Attending in person Unable to attend

Attending via teleconference
Please provide your telephone number: _____

Please reschedule so I can attend.
Proposed alternative date: (dd/mm/yy): _____ and Time: _____

Please email/fax this back to (insert email/fax number): _____

Yours sincerely (name): _____

Role: _____ Organisation: _____

This page is intentionally left blank

GP Confirmation

Organisation: _____

Palliative Care Case Conference

To:	Email/fax number:
From:	No. of pages: (including this page)
Subject: Palliative Case Conference	Date sent: (dd/mm/yy):

Dear Dr, _____

Following our recent correspondence with you a Palliative Care Case Conference has been organized for: (Resident/client name): _____

Resident/client DOB (dd/mm/yy): _____

Case conference date (dd/mm/yy): _____ Start time: _____

Expected duration: _____ Venue: _____

If you are joining by teleconference, please dial in using the following telephone number and code:

Telephone: _____ Code: _____

Reason for case conference: _____

Yours sincerely (name): _____

Role: _____ Organisation: _____

This page is intentionally left blank

Information for you and your family

Organisation:

Palliative care case conferences

It has been suggested that a case conference be held to discuss how you, or your family member might benefit from palliative care. The following explains what this is and why it is important.

Case conference: Case conferences or family meetings are an opportunity to discuss a person's care needs. They ideally include the person (if able to attend), their family and/or their substitute decision-maker, and members of the care team including the doctor.

Palliative care: Palliative care is person- and family-centred care that supports a person to live the best life they can with a life-limiting illness. A life-limiting illness means that the person has little or no prospect of cure and is expected to die. The focus is on quality of life.

Life-limiting illnesses include dementia, advanced heart, kidney, lung or liver disease, cancer, and motor neurone disease.

People can receive palliative care for days or weeks, or for months to years. Older people coming to the end of their life without illness may have some of the same care issues. They can also benefit from the approaches to care taken in palliative care.

Common care issues in palliative care include:

- pain
- dyspnoea (breathing difficulty)
- dysphagia (difficulty with swallowing)
- constipation/incontinence (bowel and/or bladder management)
- depression
- delirium (sudden confusion)
- anxiety
- nausea (feel that you want to vomit)
- fatigue (tiredness).

Who should attend a case conference?

Staff in residential aged care facilities and providers of home care often meet with families. If possible, the person receiving care should attend, their GP, and any concerned family members or friends.

Your contact for this case conference is:

Name of staff member: _____ Role: _____

Telephone: _____

This page is intentionally left blank

Invitation for you and your family

Organisation: _____

Palliative Care Case Conference

A palliative care case conference has been organised for:

Name of resident/client: _____

Resident/client date of birth (dd/mm/yy): _____

Case conference date (dd/mm/yy): _____ Start time: _____

Location: _____

Please let us know if you can attend. If you would like to join by telephone, let us know and provide a suitable number to contact you.

Your contact for this case conference is:

Name of staff member: _____

Role: _____

Telephone: _____



On the next page you will find information on palliative care and palliative care case conferences

Invitation for you and your family

Palliative Care Case Conference (continued)

Case conference: Case conferences or family meetings are an opportunity to discuss a person's care needs. They ideally include the person (if able to attend), their family and/or their substitute decision-maker, and members of the care team including the doctor.

Palliative care: Palliative care is person- and family-centred care that supports a person to live the best life they can with a life-limiting illness. This means that the person has little or no prospect of cure and is expected to die. The focus is on quality of life.

Life-limiting illnesses include dementia, advanced heart, kidney, lung or liver disease, cancer, and motor neurone disease.

People can receive palliative care for days or weeks, or for months to years. Older people coming to the end of their life without illness may have some of the same care issues. They can also benefit from the approaches to care taken in palliative care.

Common care issues in palliative care include:

- pain
- dyspnoea (breathing difficulty)
- dysphagia (difficulty with swallowing)
- constipation/incontinence (bowel and/or bladder management)
- depression
- delirium (sudden confusion)
- anxiety
- nausea (feel that you want to vomit)
- fatigue (tiredness).

Who should attend a case conference?

Staff in residential aged care facilities and providers of home care often meet with families. If possible, the person receiving care should attend, their GP, and any concerned family members or friends.

Confirmation for you and your family

Organisation: _____

Palliative Care Case Conference

A palliative care case conference has been organised for:

Name of resident/client: _____

Resident/client date of birth (dd/mm/yy): _____

Case conference date (dd/mm/yy): _____ Start time: _____

Location: _____

Your involvement in planning care is important. If you are unable to attend in person but would like to join by telephone, please dial in using the following telephone number and code.

Dial-in telephone number: _____ Code: _____

Your contact for this case conference is:

Name of staff member: _____

Role: _____

Telephone: _____

Please write down if there are any issues you want to talk about and remember to bring this form with you to the meeting so that this can be included.



On the next page you will find information on palliative care and palliative care case conferences

Confirmation for you and your family

Palliative Care Case Conference (continued)

Case conference: Case conferences or family meetings are an opportunity to discuss a person's care needs. They ideally include the person (if able to attend), their family and/or their substitute decision-maker, and members of the care team including the doctor.

Palliative care: Palliative care is person- and family-centred care that supports a person to live the best life they can with a life-limiting illness. This means that the person has little or no prospect of cure and is expected to die. The focus is on quality of life.

Life-limiting illnesses include dementia, advanced heart, kidney, lung or liver disease, cancer, and motor neurone disease.

People can receive palliative care for days or weeks, or for months to years. Older people coming to the end of their life without illness may have some of the same care issues. They can also benefit from the approaches to care taken in palliative care.

Common care issues in palliative care include:

- pain
- dyspnoea (breathing difficulty)
- dysphagia (difficulty with swallowing)
- constipation/incontinence (bowel and/or bladder management)
- depression
- delirium (sudden confusion)
- anxiety
- nausea (feel that you want to vomit)
- fatigue (tiredness).

Who should attend a case conference?

Staff in residential aged care facilities and providers of home care often meet with families. If possible, the person receiving care should attend, their GP, and any concerned family members or friends.

Staff Communication Sheet:

Organisation: _____

Palliative Care Case Conference

A palliative care case conference has been organised for:

Name of resident/client: _____

Case conference date (dd/mm/yy): _____ Start time: _____

Location: _____

As valuable members of the care team your contribution to the case conference is important. Please list below any issues, concerns or suggestions you would like mentioned. Common issues include review of symptoms (e.g. pain, dyspnoea), concerns with nutrition or hydration, family issues, emotional concerns of the resident. If you are available and would like to attend the case conference, please contact the Case Conference Facilitator:

Name of Facilitator: _____

Issue, concern or suggestion. Please be as specific as possible.	Designation

This page is intentionally left blank

Palliative Care Case Conference

Organisation: _____

Summary - Residential Care

Full name of client: _____

DOB (dd/mm/yy): _____

Purpose of Case Conference: _____

Resident consent/substitute decision-maker (SDM) consent

My care provider has explained the purpose of a case conference and I give permission for my care provider to prepare a case conference. I give permission to the providers listed below to participate in the case conference and discuss my/my family member's medical history, diagnosis, and current needs.

Signature: _____

Date: _____

Dial-in telephone number: _____ Code: _____

Resident in attendance? Yes No If no, give reason: _____

Family Members		
Name	Relationship	Attending in person (P) or teleconference (T)
		P T
		P T
		P T
		P T
		P T
Health and Care Professionals		
Name	Discipline/Position	Attending in person (P) or teleconference (T)
		P T
		P T
		P T
		P T
		P T

Palliative Care Case Conference

Summary - Residential Care (continued)

Start time: _____

Need (as appropriate): _____

Key Issues	Description
<p>Advance care plan</p> <p>Does this need to be reviewed? Does the person understand their diagnosis/prognosis?</p>	
<p>Symptoms</p> <p>For example: fatigue, anorexia, pain, nausea, dyspnoea, dysphagia</p>	
<p>Social/psychological needs</p> <p>For example: isolation, anxiety, depression What supports are being provided? What supports are needed?</p>	
<p>Assessments/investigations</p> <p>Can the resident manage ADL's (Activities of Daily Living)? Do they need additional support?</p>	
<p>Carer/Family issues or needs</p>	
<p>Other</p> <p>For example: general issues, housing issues, financial issues</p>	

Palliative Care Case Conference

Summary - Residential Care (continued)

Agreed Action Plan

Goal	Actions	Key Person(s) Responsible	Description

Palliative Care Case Conference

Summary - Residential Care (continued)

Time completed: _____

General Practitioner: _____

Tick appropriate box

Original placed in the **resident's** clinical notes

Copy provided to all participants

Copy sent to GP

Resident's care plan and assessment reviewed and updated

Palliative Care Case Conference Facilitator

Name: _____ Position: _____

Signature: _____ Date (dd/mm/yy): _____

Medicines from the PBS Prescriber bag for terminal phase symptoms

PBS Item Code	Pharmaceutical benefit and form	Strength	Packet size	Max qty (packs)	Max qty (units)
3451P	Adrenaline (Epinephrine) injection	1 in 1000 (1 mg/mL)	5 x 1mL amps	1	5
3478C	Clonazepam oral liquid	2.5 mg/mL (0.1 mg/drop)	1 x 10mL	1	1
3466K	Furosemide (Frusemide) ampoule	20 mg/ 2 mL	5 x 2mL	1	5
3456X	Haloperidol ampoule	5 mg/mL	10 x 1mL	1	10
3470P	Hydrocortisone Sodium Succinate injection* OR	100 mg (reconstituted to 2mL) OR	Single injection	2	2
3471Q	Hydrocortisone Sodium Succinate injection*	250 mg (reconstituted to 2mL)	Single injection	1	1
3473T	Hyoscine Butylbromide ampoule	20 mg/mL	5 x 1mL	1	5
3476Y	Metoclopramide ampoule	10 mg/ 2 mL	10 x 2mL	1	10
10178Q	Midazolam ampoule	5 mg/mL	10 x 1mL	1	10
10862Q	Morphine ampoule OR	10 mg/mL OR	5 x 1mL	1	5
3479D	Morphine ampoule OR	15 mg/mL OR	5 x 1mL	1	5
10868B	Morphine ampoule OR	20 mg/mL OR	5 x 1mL	1	5
3480E	Morphine ampoule	30 mg/mL	5 x 1mL	1	5
10786Q	Naloxone hydrochloride injection OR	400 microgram/mL OR	5 x 1 mL	2	10
11233F	Naloxone hydrochloride injection	400 microgram/mL	10 x 1 mL	1	10

Based on the emergency practice concept proposed by Seidel et al 2006 Aust Fam Physician. 2006 Apr;35(4):225-31. Information from PBS listings current as of March 2024. See www.pbs.gov.au for more.

Medicines from the PBS Prescriber bag for terminal phase symptoms

Many people with palliative care needs, choose to be cared for and die at home.

Prescribing medicines in advance (anticipatory prescribing), ensures prompt response when symptoms occur. Yet, people can deteriorate suddenly and rapidly.

In Australia, some medicines are provided without charge to prescribers, who can supply them free on home visits. The Pharmaceutical Benefits Scheme (PBS) prescriber bag list includes medicines which can be useful in caring for the dying, in the home environment. These can:

- Be administered immediately, to manage symptoms; and
- See the person through until a prescription can be dispensed.

A prescriber bag supply order form can be ordered online from [Services Australia](#). The forms allow monthly ordering of medicines. They must be completed, signed, and given to a community pharmacist for dispensing.

The PBS prescriber bag is a safety net for those who deteriorate suddenly at the end of life. It is not a substitute for good advanced planning.

Symptoms common in the terminal phase

Agitation, or terminal restlessness:

Characterised by anguish (spiritual, emotional, or physical), restlessness, anxiety, agitation, and cognitive failure. Sublingual clonazepam, subcutaneous midazolam and subcutaneous haloperidol may be used.

Delirium: Haloperidol is commonly used to reduce distress due to delirium. For severe hyperactive delirium with agitation see response to agitation.

Dyspnoea: Subcutaneous morphine is the gold standard. Avoid repeated dosing in people with serious kidney failure. Because there may be an anxiety

component, sublingual clonazepam or subcutaneous midazolam may also have a role. Nebulised adrenaline may give temporary relief if stridor is present.

Nausea and vomiting: For onset of new nausea or vomiting in the terminal phase, when the cause is unknown, haloperidol or metoclopramide are usually used as first-line therapy.

Oedema associated with heart failure: Intravenous or subcutaneous furosemide (frusemide) can be adjusted against the oral dose until symptoms are controlled.

Pain: Subcutaneous morphine can be used in most people with pain. Avoid repeated dosing in severe renal failure. Naloxone can be used for opioid poisoning.

Respiratory tract secretions: The inability to clear secretions from the oropharynx or trachea causes pooling of fluids in the throat. This results in rattly breathing. This may be more distressing for the people around than for the person themselves. Hyoscine butylbromide can be used.

Rigidity associated with end-stage Parkinson disease: If dopaminergic medication is ceased, subcutaneous midazolam or sublingual clonazepam may help to relieve rigidity.

Palliative care emergencies: A sudden and life-threatening change in a person's condition, may be unexpected. Some can be foreseen, based on the nature and location of the disease. These may include:

- Superior vena cava obstruction;
- Catastrophic haemorrhage;
- Airway obstruction;
- Seizures; and
- Spinal cord compression.

Morphine and midazolam, administered subcutaneously in the home, can reduce distress. Subcutaneous hydrocortisone may be used in place of

dexamethasone, where an anti-inflammatory is useful (e.g. bowel obstruction, spinal cord suppression, and airway obstruction).

*Note: The final volume of the hydrocortisone, once reconstituted, is 2mL which may limit the dose that can be comfortably administered.

In managing a bowel obstruction, hyoscine butylbromide is helpful in managing the cramping pain, while haloperidol is preferred for nausea and vomiting. Topical use of adrenaline is suggested for small volume superficial malignant bleeding.

Practical tips

In addition to carrying medicines, bring equipment to administer them.

Order your PBS prescriber bag medicines at the end of the month.

Lock medicines up in a secure place – you are responsible for their security and must adhere to legislative requirements around secure storage of S8 medications such as opiates.

Keep a recording book for administering, supplying, or discarding medications.

The maximum volume generally accepted for a subcutaneous injection is around 1.5mL - larger volumes are more likely to be associated with pain at the injection site. See hydrocortisone and volume considerations above.

Speak with the community pharmacist about stocking these medicines, so the ongoing prescription can be dispensed immediately.

For specific dosing advice, refer to:

- [Australian Medicines Handbook](#)
- [Palliative Care Therapeutic Guidelines](#)
- palliAGED [symptoms and medicines](#)
- [CareSearchgp app](#).

MBS Items for Nurse Practitioners

Note: This webpage aims to provide clear guidance on the remuneration that Nurse Practitioners can obtain for palliative care services for patients within the community.

MBS	Medicare Initiative	MBS Benefit 85% (as of 01/11/2023)
82200	Professional attendance by a participating nurse practitioner for an obvious problem characterised by the straightforward nature of the task that requires a short patient history and, if required, limited examination and management.	\$9.00
82205	Professional attendance by a participating nurse practitioner lasting less than 20 minutes and including any of the following: taking a history; undertaking clinical examination; arranging any necessary investigation; implementing a management plan; providing appropriate preventive health care, for 1 or more health related issues, with appropriate documentation.	\$19.60
82210	Professional attendance by a participating nurse practitioner lasting at least 20 minutes and including any of the following: taking a detailed history; undertaking clinical examination; arranging any necessary investigation; implementing a management plan; providing appropriate preventive health care, for 1 or more health related issues, with appropriate documentation.	\$37.15
82215	Professional attendance by a participating nurse practitioner lasting at least 40 minutes and including any of the following: taking an extensive history; undertaking clinical examination; arranging any necessary investigation; implementing a management plan; providing appropriate preventive health care, for 1 or more health related issues, with appropriate documentation.	\$54.85

There are also a number of [MBS Telehealth items](#) (video and phone services) for a range of (out of hospital) consultations.

Source: <https://www.servicesaustralia.gov.au/bulk-billing-for-nurse-practitioners-and-midwives>. Accessed 16/01/2024

This page is intentionally left blank

MBS Remuneration to Support a Planned General Practice Palliative Care Pathway (Residential Aged Care)

The aim of this webpage is to support multidisciplinary practice by describing a palliative care pathway which clearly shows remuneration for palliative care services.

Suggested Timeframe	Medicare Initiative	Activities	MBS Item	MBS Benefit 100% (as of 01/11/23)
0 months	Comprehensive Medical Assessment (CMA)	On admission, then annually. Identify who is appointed to assist with healthcare decisions for patients who do not have 'capacity' for palliative care discussions. Select relevant item based on complexity and PN +GP time.	701	\$65.30
			703	\$151.80
			705	\$209.45
			707	\$295.90
	Domiciliary Medication Management Review (HMR)	Referral to eligible pharmacist; ensures optimal management of patient with 5 or more medications and/or complexity.	903	\$116.70
2 week	GP Mental Health Treatment Plan	Select relevant item depending on time and GP training.	2700	\$78.95
			2701	\$116.20
			2712	\$78.95
			2713	\$78.95
			2715	\$100.20
			2717	\$147.65
1st month	Care Plan Contribution	For patients with chronic disease; GP contributes to facility's plan. GP contribution entitles the patient to Medicare allied health services (EPC), 5 per calendar year.	731	\$77.50
2nd month	Case Conference	Opportunity for holistic informed approach to ongoing care for providers, carers, and family. Organised by the GP; 20-40 minutes long; requires GP and at least 2 other providers (e.g. Palliative Care Specialist) in 'real' time.	739	\$133.10

Suggested Timeframe	Medicare Initiative	Activities	MBS Item	MBS Benefit 100% (as of 01/11/23)
4th month	Level D consultation	To complete Advance Care Directive, following earlier discussions.	44	\$117.40
5th month	Review GP Mental Health Treatment Plan	4 weeks - 6 months after preparation of plan, review referral feedback and progress against goals.	2712	\$78.95
6 month	Care Plan Contribution	Review of facility's multidisciplinary plan and above GP contribution.	731	\$77.50
8 month	Case Conference	Organised by GP; 20-40 minutes long; GP+2 other providers in 'real' time.	739	\$133.10
After 12 months	Repeat CMA, case conferences and care plan contributions where clinically required.		As above.	

Source: Based on information from: PHN North Western Melbourne, MBS Remuneration to Support Planned End-of-Life and Palliative Care for Patients, A guide for Health Professionals working in General Practice and Residential Aged Care. Sept. 2019.